

Breast Preop instructions Instructions

Pre-op Clinic Plan Instructions

Use the following smartsets:

Surgical Oncology Day of Surgery Orders

SurgOnc Clinic Preop

Place the following under SurgOnc Clinic Preop

1. OR Case request (see options listed for procedure; choose appropriate one)

<input type="checkbox"/>	LUMPECTOMY, BREAST	RUHLMPECTMY	General
<input type="checkbox"/>	LUMPECTOMY, BREAST, WITH NEEDLE LOCALIZATION	RUHLMPECTMYNDL	General
<input type="checkbox"/>	MASTECTOMY, MODIFIED RADICAL	MASTMODR	General
<input type="checkbox"/>	MASTECTOMY, MODIFIED RADICAL, WITH SENTINEL AND AXILLA...	RUHMASTCMY	General
<input type="checkbox"/>	MASTECTOMY, PARTIAL	LUMPBR	General
<input type="checkbox"/>	MASTECTOMY PARTIAL/LUMPECTOMY MAGSEED LOCALIZATION	LUMPAGSE	General
<input type="checkbox"/>	MASTECTOMY PARTIAL / LUMPECTOMY NEEDLE LOCALIZATION	LUMPNDL	General
<input type="checkbox"/>	MASTECTOMY, RADICAL	MASTRAD	General
<input type="checkbox"/>	MASTECTOMY, ROBOT ASSISTED	MASTROBO	General
<input type="checkbox"/>	MASTECTOMY, SIMPLE	MASTSIMP	General
<input type="checkbox"/>	MASTECTOMY, SKIN SPARING OR NIPPLE SPARING	MASNIPSP	General

Under the Breast/Melanoma Section under Radiology, place the proper orders needed

1. For partial mastectomy/lumpectomy/breast excisional biopsy cases, breast localization is generally needed (unless lesion is palpable and attending instructs that localization is not needed)

2. If **Magseed** localization is needed, order the following:

- US or Mammo Guided Magseed placement (depends on which modality was used to biopsy the breast, US or mammo)
- Mammo Clip Placement (LT or RT)
- XR Surgical Specimen (to check the magseed in specimen)
- US of surgical specimen (only if seed was placed by US)

3. If **Needle** localization is needed, order the following

- US or Mammo guided Needle localization (specify laterality)
- XR surgical specimen if needle placed by mammo
- US surgical specimen if needle placed by US

4. For identifying Sentinel LN

- Choose NM Sentinel Node Breast Injection (unless attending says otherwise; IE MAGTRACE**)- **If using magseed or magtrace, make sure this is indicated in special needs in the OR case request.**
- Discuss with attending in patients undergoing neoadjuvant chemotherapy for node positive disease if magseed localization is needed as well for the axilla to identify the LN

▼ Radiology

- ▶ Xray
- ▶ Cardiac Studies
- ▼ Breast/Melanoma
 - Mammo Clip Placement LT ■
STAT
 - Mammo Clip Placement RT ■
STAT
 - Mammo Guided Magseed Placement Left ■
STAT
 - Mammo Guided Magseed Placement Right ■
STAT
 - US Guided Magseed Placement Left ■
STAT
 - US Guided Magseed Placement Right ■
STAT
 - US Guided Breast Needle Localization ■
 - Mammo Guided Breast Needle Localization ■
 - XR Surgical Specimen Radiology ■
 - US Surgical Specimen - Ultrasound ■
 - NM Sentinel Node Breast Injection Bilateral - No Imaging ■
 - NM Sentinel Node Breast Injection Left - No Imaging ■
 - NM Sentinel Node Breast Injection Right - No Imaging ■
 - NM Lymphatic Scan ■

Order the indicated blood tests (cbc, bmp), cxray, EKG's from the order set





Place the following under Consults if needed:

1. Ambulatory referral to Stop Smoking Program
 - Patients should stop smoking and drinking 4 weeks prior to surgery
2. Ambulatory referral to Diabetic education or Endocrinology for assistance in diabetic control if poorly controlled DM



Open the Surgical Oncology Day of Surgery smartest

- Select Pre-op Cleaning
- Select no venipuncture and select arm that is not on the surgery side (If applicable)
- Select multi-modal meds, pick meds appropriate for patient

These orders are intended for Day of Surgery only

▼ GENERAL ORDERS

▶ NPO Click for more

▼ Surgical/Pre-operative Cleaning Preparation

- Surgical / Procedure Preparation: Pre-op Cleaning
 Until discontinued, Starting 2/25/21, Please clean patient's abdomen with Chlorhexidine Gluconate 2% solution (Dyna-Hex) for 30 seconds and allow to dry prior to the patient going to OR. If the patient is allergic to Chlorhexidine, notify Provider on-call and use other antiseptic solution for pre-op., Pre-op

▼ BREAST

▼ Nursing - No IV On Surgery Side

- No Venipuncture
 Until discontinued, Starting 2/25/21, No IV on side of surgery, Pre-op

▼ Medications

- acetaminophen (TYLENOL) tablet
 1,000 mg, Oral, Once, Pre-op
- gabapentin (NEURONTIN) capsule
 600 mg, Oral, Once, Pre-op
- gabapentin (NEURONTIN) capsule (if >70yo or CrCl <30)
 300 mg, Oral, Once, Pre-op
- scopolamine (TRANSDERM-SCOP) 1.5 mg patch
 1 patch, Transdermal, Apply one patch behind ear and discontinue 24 hours after surgery, Pre-op

Melanoma Preop instructions Instructions

1. Follow the same instruction using the SurgOnc Clinic Preop and SurgOnc Day of Surgery orders
2. Under the Breast/Melanoma Section in the SurgOnc Clinic Preop select NM Lymphatic Scan if patient needs sentinel LN biopsy

